




# IHDP Growth Percentiles: LBW Premature Girls<sup>1,2</sup>



Growth of very-low-birth-weight (VLBW,  $\leq 1500$  g) and low-birth-weight (LBW, 1501 to 2500 g) premature ( $\leq 37$  weeks, GA) infants differs from that of normal-birth-weight term infants during infancy and early childhood. Because these infants may not catch up to term infants in growth during the early years, their growth should be compared to that of premature infants of similar birth weight.

The growth percentiles presented here are based on a large sample of infants enrolled in the Infant Health and Development Program (IHDP).<sup>1,2</sup> Some infants most likely to experience growth problems from biologic or environmental causes, premature infants with birth weight greater than 2500 g, and small-for-gestational-age term infants were excluded.<sup>1</sup> Study infants, however, are probably typical of premature infants who receive modern neonatal intensive care.

## References

1. The Infant Health and Development Program: Enhancing the outcomes of low-birth-weight, premature infants. *JAMA* 1990;263(22):3035-3042.
2. Casey PH, Kraemer HC, Bernbaum J, et al: Growth status and growth rates of a varied sample of low birth weight, preterm infants: A longitudinal cohort from birth to three years of age. *J Pediatr* 1991;119:599-605.

IHDP studies were supported by grants from the Robert Wood Johnson Foundation, Pew Charitable Trusts, and the Bureau of Maternal and Child Health, US Department of Health and Human Services. These graphs were prepared by SS Guo and AF Roche, Wright State University, Yellow Springs, Ohio. IHDP, its sponsors, and the investigators do not endorse specific products.

## Instructions for Use

1. Measure and record weight, length, and head circumference.
2. Calculate gestation-adjusted age by subtracting Adjustment for Prematurity in weeks from postnatal age in weeks. Adjustment for Prematurity equals 40 weeks minus GA. For example, at 12 wk postnatal age, an infant born at 30 wk GA would be 2 wk (0.5 mo) gestation-adjusted age.
3. Plot data at the gestation-adjusted age on the appropriate graph.
4. When possible, plot serial data on the same graph to permit detection of change in growth percentiles with age.

## Interpretation

These graphs permit comparison of a LBW premature girl's growth relative to current reference data. Further investigation may be indicated when the plotted measurements are markedly different from the 50th percentile, or growth percentile changes rapidly.

Provided as a service of  
**SIMILAC® NeoCare™**  
Premature Infant Formula With Iron